

Fig. 1.




Fig. 2.

1049-XNP-2

Wick Time & Tensile vs. Cantilever Stiffness

FIGURE 3

PAGE 3




FIGURE 4

Fluid Transfer to core by time

PAGE 4





FIG. 5

Acquisition of Fluid of 4th 75ml Insult
Using Saddle Wicking Apparatus



Procedure = 75ml insults
every 20min

FIG. 6
PAGE 6




FIG. 7
PAGE 7
Distribution of Fluid within Training Pant Using Saddle Wicking Apparatus




FIG. 8

PAGE 8

Distribution of Fluid within Training Pant Using Saddle Wicking Apparatus




FIG. 9
PAGE 9

**Acquisition of Fluid of 3rd 75ml Insult
Using Market Pulp Flat Acquisition Test**






FIG. 10
PAGE 10

Acquisition Rate of Training Pants
Using Market Pulp Flat Acquisition Test



Acquisition of Fluid of 4th 75ml Insult
Using Market Pulp Flat Acquisition Test



Procedure = 75ml insults
every 20min

10021658.120701




Fig. 12A.




Fig. 12B.




Fig. 12C.

FIG. 12A, 12B, 12C




Fig. 13A.




Fig. 13B.




Fig. 13C.




Fig. 13D.

10021568.120701